

L305 ANSWER 91 OF 145 TOXCENTER COPYRIGHT 2003 ACS

AN 1987:154263 TOXCENTER

CP Copyright 2003 ACS

DN CA10725229106Q

TI Free radical and cytotoxic effects of **chelators** and their iron **complexes** in the hepatocyte

AU Mostert, L. J.; Van Dorst, J. A. L. M.; Koster, J. F.; Van Eijk, H. G.; Kontoghiorghes, G. J.

CS Dep. Chem. Pathol. Biochem. I, Erasmus Univ., Rotterdam, Neth..

SO Free Radical Research Communications, (1987) Vol. 3, No. 6, pp. 379-88.
CODEN: FRRCEX. ISSN: 8755-0199.

CY NETHERLANDS

DT Journal

FS CAPLUS

OS CAPLUS 1987:629106

LA English

ED Entered STN: 20011116

Last Updated on STN: 20021105

AB In a comparative screening study of **chelators** intended for clin. use eleven iron **chelators** have been tested for their ability to mobilize ⁵⁹Fe from ⁵⁹Fe-labeled ferritin and from hepatocytes of rats labeled with ⁵⁹Fe-transferrin. The toxic effects of the **chelators** were also studied using microsomal lipid peroxidn. induced by Fe³⁺/ADP and NADPH. From these tests it was shown that 1,2-di-Me 3-hydroxypyrid-4-one (L1) and mimosine were the most effective Fe **chelators** in Fe mobilization and did not catalyze lipid peroxidn. Thus, aside from their Fe binding properties, **chelators** should be examd. for their role in catalyzing lipid peroxidn. in toxicol. screening.